

Shower spectrum and detection of tau from Earth-skimming neutrinos

Ming-Huey A. Huang ^{a *}

^aGeneral Education Center, National United University
1 Lien-Da, Kung-Ching Li, Miao-Li, TAIWAN. 36003, ROC

A realistic Monte-Carlo simulation code is developed to study the shower spectrum initiated by decay of tau leptons, which come from charged current interaction inside the Earth. AGN and GZK neutrino fluxes are used to simulate the shower spectrum.

At neutrino energy above 10^{15} eV , detecting neutrinos skimming through the Earth provides an unique chance to detect those high energy neutrinos and confirmation of $\nu_\mu \rightarrow \nu_\tau$ oscillation. To evaluate the detectable event rate and optimize the detection efficiency, a realistic Monte-Carlo simulation code is developed.

The Earth is modeled as a spherical sphere of several layers of materials and density. The $\nu - N$ cross-section is modeled by CTEQ-6 model [1]. For this study, the energy loss of τ is modeled by continuous energy loss, i.e.,

$$-dE/(\rho dx) = \alpha + \beta E = \beta E,$$

where ρ is density of material, dx is distance traveled by τ , and α and β is the coefficients of energy loss from all processes [1]. The τ survival probability P is modeled by

$$dP/P = -dx/(\kappa E) = dE/(\kappa \beta \rho E^2),$$

where κE is decay length of τ at energy E in PeV and $\kappa = 49.9 \text{ m/PeV}$.

τ decay is simulated with TAUOLA package with fully polarized τ s. When τ s decay inside the Earth, the ν_τ s are re-propagated inside the Earth and could generate new τ s again. When a τ decays outside the Earth, it initiate air shower, except when τ decay to muon. The shower energy is sum of the energy from electrons and hadrons. The shower simulation is done by a modified version of CORSIKA [2].

*This study is supported by National United University and grant NSC-92-2119-M-239-001 from National Science Council of Taiwan, ROC.

To compare with previous results from analytical method [1], ν_τ s are injected to a fix width of 100km of rock. Shown in the Fig. 1, the τ spectrum from both methods agrees quite well. This test conclude that the algorithm used in this Monte-Carlo simulation is correct. We are updating energy loss with Stochastic processes and the complete study will be published soon.




Figure 1. Spectrum of τ from ν_τ pass through 100 km of rock from AGN (left figure) [3] and GZK (right figure) [4]. The blue markers come from analytical calculation [1], while the red markers come from Monte-Carlo simulation in this study.

REFERENCES

1. J.J. Tseng et al., Phys. Rev. **D68**, (2003),063003 and references therein.
2. M.A. Huang, C.C. Hsu, and P. Yeh, *Proc. of the 28th Int. Cosmic Ray Conf.*, Tsukuba, Japan, (2003), 589.
3. O.E. Kalashev, et al., Phys. Rev. **D66**, (2002), 063004.
4. R. Engel, D. Stecker, and T. Stanev, Phys. Rev. **64**, (2001), 093010.